

✓ Distribution and Population Constitution of
Drosophila in South East Asia and Oceania, IV.

New and unrecorded species of Drosophilidae
in Solomon Islands

Haruo Takada

Department of General Education, Sapporo University

謹呈岡田曲登日先生

Reprinted from the Journal of the Faculty
of General Education, Sapporo University

No. 22: 139~145

Sapporo, Japan
March, 1983

著者

Distribution and Population Constitution of *Drosophila* in South East Asia and Oceania, IV.

New and unrecorded species of *Drosophilidae*
in Solomon Islands

by

Haruo Takada

Department of General Education, Sapporo University

Collecting trips were twice in Southeast Asia and Oceania from June 25 to August 31 in 1972, and June 22 to August 31 in 1973, to explore the drosophilid flies. A lot of specimens of drosophilid flies were collected by net-sweeping and exposing fermenting banana baits from two tropical islands, Guadalcanal and New Georgia, Solomon Islands, from August 1 to August 9, 1973 with Dr. Eizi Momma.

They represent twelve species belonging to the *Drosophilidae* with five genera including a new species in subgenus *Hirtodrosophila*, genus *Drosophila*. These new flies were captured by a sweeping-net over on small brownish fungi which grew on some fallen trees at the Honiara Rifle Range, Honiara, Guadalcanal.

The author's hearty thanks are due to Dr. Eizi Momma for giving the public expenses from research grant of the Scientific Research Found of the Ministry of Education of Japan in 1973.

All type specimens are deposited at the Biological Laboratory, Sapporo University.

Haruo Takada

Family Drosophilidae

Subfamily Steganinae

Genus *Leucophenga* Mik

Subgenus *Leucophenga* Mik

- ✓ 1) *Leucophenga gibbosa* (de Meijere), 1914, Tijdschr. Ent. 57 : 264.
Specimen examined: ♂, Aug. 6, 1973, Munda, New Georgia
(Takada).

Subfamily Drosophilidae

Genus *Drosophila* Fallén

Subgenus *Drosophila* Fallén

- ✓ 2) *Drosophila pararubida* Mather, 1961, Univ. Qd Pap. 1 : 251.
Specimens examined: 6 ♂♂, 7 ♀♀, Guadalcanal, Aug 2,
1973 and 2, New Georgia, Aug. 6, 1973 (Takada).

syn. of
hypocausta

- ✓ 3) *Drosophila quadrilineata* de Meijere, 1911, Tijdschr. Ent.
54 : 396.
Specimens examined : 2 ♂♂, Guadalcanal, Aug. 4, 1973
(Takada).

Subgenus *Hirtodrosophila* Duda

- ✓ 4) *Drosophila menisigra* Bächli, 1973, Mitt. Zool. Berl., 49 : 309.
Specimens examined: 2 ♂, 2 ♀ Guadalcanal, Aug. 3,
1973 (Takada).
- ✓ 5) *Drosophila seminigra* Duda, 1926. Suppl. Ent. 14 : 68.
Specimens Examined.: 2 ♂♂, Guadalcanal, Aug. 3, 1973
(Takada).
- ✓ 6) *Drosophila solomonica* Takada n. sp. (Fig. 1)
D. (Hirtodrosophila) solomonica ^{nom. nud.} n. sp., Takada 1979. Zool.
Mag., 88, 4 : 673, at the 50th Annual Meeting, Zool Soc. of
Japan (in Japanese).




Fig. 1 *Drosophila* (*Hirtodrosophila*) *solomonica* Takada, n. sp.
1. Phallic organ. 2. Periphallal organ. 3. Caudal margin of epandrium with
semitransparent tape-like process. 4. Spermatheca. 5. Egg-guide

Haruo Takada

Male & Female. Body about 1.0-1.3 mm, wing length about 1.4 mm, small species. Front, mesonotum, scutellum and humerus coffee brown, front somewhat velvety. Eye deep purplished red with piles. The third joint of antenna with grey and long hairs anteriorly. Arista with three dorsal and one ventral branches besides a large fork. Ocellar triangle brown. Anterior reclinate orbital bristle one-third posterior reclinate orbitals, half of proclinate bristle. Proclinate orbitals inside others, anterior reclinate nearer to proclinate than to posterior reclinate. Carina high, broad. Cheeks yellow. Palpus yellow with a apical stout seta. Vibrissa long and black, other orals fine. Acros-tichal hairs in six rows. Thoracic pleura, abdomen, legs and halteres are yellow.

Wings hyaline, R2 + 3 straight and short, R4 + 5, and M parallel; Costal-index about 0.8; 4 V-index about 2.6; 4 C-index about 2.3; 5 x-index about 3.0 and AC-index about 4.0. Cl-bristles two, subequal. C3-fringe on basal half.

Periphallic organs: Pale yellow. Epandrium pubescent, truncate below, posterior margin with few bristles, narrowly elongated below at heel, toe rounded with two stout bristles, without invagination at the insertion of surstylus. Surstylus broad, crescent, distally concaved and with about seven pointed black teeth, caudal margin of epandrium *with about seven semitransparent tape-like bristles*. Cercus oval, pubescent and hairy, *undermargin with thmb-shaped process*.

Phallic organs: Pale yellow. Adeagus pale yellow, slender, apically swollen and membranous. Apodeme of aedeagus pale yellow, narrow and short. Anterior paramere elongate, apparently fused to ventral fragma, with a few apical sensilla. Hy-

Distribution and Population Constitution of *Drosophila* in
South East Asia and Oceania IV.

pandrium small and rounded, with submedian spine. Ventral fragma V-shaped without median process. Phallosomal index about 3.0.

Egg-gude: Pale yellow, spindle-shaped, apically tridently, ending on a black stout tooth, respectively. Basal isthmus long and broad.

Holotype: Male, The Honiara Rifle Range, Honiara, Guadalcanal, Solomon Is., Aug. 3, 1973 (Takada).

Allotype: Female, collected together with the holotype.

Paratypes: 2 males and 2 female, collected together with the holotype.

Collecting site: Small brownish fungi on a fallen tree. Same fungi also attracted two other species of *Hirtodrosophila*, *D. (H.) seminigra* Duda, and *D. (H.) menisigra* Bächli, and with *Mycodrosophila amabilis* (de Meijere).

Relationships: Closely allied to *D. (H.) seminigra* Duda, 1926, but differs from it in shapes of phallic and periphallic organs and egg-guides (Bächli, 1973).

Slide No. of genitalia: SGHZ-5, -6, -9 (Sapporo University).

Subgenus *Sophophora* Sturtevant

✓ (7) *Drosophila pallidosa* Bock & Wheeler, 1972, Univ. Tex. Publ., 7213: 38.

Specimens examined: Many, Aug. 2-4, 1973, Guadalcanal (Takada), and Aug. 5-9, 1973, New Georgia (Takada).

Front rather broad, ocellus ruby colors and abdomen yellow.

8) *Drosophila pseudoananassae* Bock, 1971, Univ. Tex. Publ., 7103: 274.

Specimens examined: Many, Aug. 2-4, 1973, Guadalcanal (Takada) and Aug. 5-9, 1973, New Georgia (Takada).

Genus *Lissocephala* Malloch

✓ 9) *Lissocephala metallescens* (de Meijere), 1914, Tijdschr. Ent. 57: 265.
already recorded from Solomon
by W. H. S. 1964

Specimen examined: ♂, Aug. 2, 1973, Guadalcanal (Takada).

Genus *Mycodrosophila* Oldenberg

✓ 10) *Mycodrosophila amabilis* (de Meijere), 1911, Tijdschr. Ent. 54: 405.

Specimens examined: 6 ♂, 4 ♂, Guadalcanal, Aug. 3, 1973 (Takada).

✓ 11) *Mycodrosophila gressitti* Wheeler & Takada, 1964. Insect of Micronesia, 14, 6: 204.

Specimens examined: ♂ ♀, Guadalcanal, Aug. 3, 1973 (Takada). Body length about 1.5 mm.

Genus *Sphaerogastrella* Duta

✓ 12) *Sphaerogastrella javana* (de Meijere), 1911, Tijdschr. Ent. 54: 422. (Fig. 2)

Specimens examined: ♂, Aug. 2, 1973, Guadalcanal (Takada) and 4 ♀, Aug. 5, 1973, New Georgia (Takada).

Distribution and Population Constitution of *Drosophila* in
South East Asia and Oceania IV.




Fig. 2 *Sphaerogastrella javana* (de Meijere)
1. Periphallic organs. 2. Phallic organs. 3. Ejaculatory
apodeme. 4. Egg-guide. 5. spermatheca.

2139

表題 - New and unrecorded species 及び文中に "unrecorded" の

附註あり。

2140 *D. pararubida* (3 *D. hypocausti* の synonym. [Whelan, 1981:42
hypocausti (3 Solomon Is.) に記述あり。]

2144 *Lissocephala metallescens* 3 Solomon Is.) に記述あり。]