

Paik, Y. K., and Kim, K. W.
Local key to species of Drosophilidae collected so far in South Korea.

DIS 31: 153-154 (1957)

collection localities have been increased, we have attempted to survey the distribution patterns of those species, as summarized below. Roman numerals indicate (I) Mt. Moodung (located in Kwangju, Chunnam province), (II) Mt. Chiri (in Kurae, Chunnam province), (III) Mt. Hanra (in Chaeju, Quilpart Island), (IV) Mt. Sori (in Kwangnung, Kyongi province), and (V) Mt. Taepaik (in Hwangjee, Kangwon province).

Species	Localities				
	I	II	III	IV	V
<i>Amiota alboguttata</i>	-	+	-	-	-
<i>A. variegata</i>	+	+	-	+	+
<i>Leucophenga argentosa</i>	+	-	-	-	+
<i>L. magnipalpis</i>	+	-	-	-	-
<i>L. concilia</i>	+	-	-	-	-
<i>L. maculata</i>	+	-	-	-	-
<i>L. quinquemaculipennia</i>	-	+	-	-	-
<i>L. ornatipennis</i>	+	-	-	-	-
<i>Mycodrosophila poecilogastra</i>	+	-	-	-	-
<i>Mycodrosophila</i> sp.	+	-	-	-	-
<i>Microdrosophila</i> sp.-1	-	+	+	-	-
<i>Microdrosophila congesta</i>	-	+	-	-	+
<i>Microdrosophila</i> sp-2	-	-	+	-	-
<i>Scaptomyza disticha</i>	+	+	+	-	+
<i>S. graminum</i>	-	+	+	+	+
<i>S. polygonia</i>	-	-	-	-	+
<i>Drosophila</i> (H) <i>alboralis</i>	-	+	+	+	-
<i>D.</i> (H) <i>sexvittata</i>	+	-	+	+	+
<i>D.</i> (H) <i>quadrivittata</i>	-	+	-	-	-
<i>D.</i> (H) <i>nokogiri</i>	-	-	+	+	-
* <i>D.</i> sp. of <i>Hirtodrosophila</i>	-	+	-	+	-
<i>D.</i> sp. close to <i>histrio</i>	-	-	+	+	+
<i>D.</i> (<i>Dorsilopha</i>) <i>busckii</i>	+	-	-	-	-
<i>D.</i> (<i>Pholadoris</i>) <i>coracina</i>	+	+	-	+	+
<i>D.</i> (<i>Pholadoris</i>) <i>puncticeps</i>	+	-	-	-	-
<i>D.</i> (<i>Pholadoris</i>) <i>rubifrons</i>	+	-	-	-	-
<i>D.</i> (<i>Sophophora</i>) <i>suzukii</i>	+	+	+	+	+
<i>D.</i> (S) <i>takahashii</i>	+	-	-	-	-
<i>D.</i> (S) <i>lutea</i>	+	-	+	-	-
<i>D.</i> (S) <i>melanogaster</i>	+	-	+	+	-
<i>D.</i> (S) <i>magnipectinata</i>	-	-	+	-	-
<i>D.</i> (S) <i>auraria</i>	+	+	+	+	+

<u>Species</u>	<u>Localities</u>				
	I	II	III	IV	V
D. (S) <i>bifasciata</i>	+	-	+	-	+
D. (S) sp. of <i>obscura</i> group	-	-	+	-	+
D. (S) <i>helvetica</i>	+	-	-	-	-
D. (Drosophila) <i>transversa</i> complex	+	+	+	+	+
D. (D) <i>nigromaculata</i>	+	-	-	-	+
D. (D) sp. close to <i>kuntzei</i>	-	-	-	-	+
D. (D) <i>testacea</i>	-	-	-	+	+
D. (D) <i>bizonata</i>	+	+	+	+	+
D. (D) <i>makinoi</i>	-	-	-	-	+
D. (D) <i>immigrans</i>	+	+	+	-	-
D. (D) sp. of <i>immigrans</i> group	-	-	+	-	-
D. (D) <i>virilis</i>	-	-	+	+	-
D. (D) <i>sordidula</i>	-	-	-	+	-
D. (D) <i>lacertosa</i>	+	-	+	+	+
D. (D) <i>cheda</i>	+	+	-	+	-
D. (D) <i>repleta</i>	-	-	+	-	-
D. (D) <i>histrio</i>	+	+	+	+	+
D. (D) sp. of <i>quinaria</i> section	+	+	+	+	-
**D. (D) sp. <i>-sternopleuralis</i>	+	-	-	-	-
D. sp. of <i>Drosophila</i>	+	-	-	-	-

*D. sp. of *Hirtodrosophila* is recorded in Okada's MS as D. histrioides.

**D. (D) sp. is recorded in Okada's MS as D. sternopleuralis.